

AD-A163 625

A CONVERGENCE THEOREM FOR NEWTON'S METHOD IN BANACH
SPACES(U) WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH
CENTER T YAMAMOTO OCT 85 ARC-TSR-2879 DRAG29-80-C-0041

1/1

UNCLASSIFIED

F/G 12/1


NL

END

FILMED

1-4

57C



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

2

MRC Technical Summary Report #2879

A CONVERGENCE THEOREM FOR NEWTON'S
METHOD IN BANACH SPACES

Tetsuro Yamamoto

AD-A163 625

**Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705**

October 1985

(Received September 17, 1985)

DTIC
ELECTE
FEB 5 1986
S B D

DTIC FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

86 2 5

036

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

A CONVERGENCE THEOREM FOR NEWTON'S
METHOD IN BANACH SPACES

Tetsuro Yamamoto*

Technical Summary Report #2879
October 1985

ABSTRACT

On the basis of the results obtained in a series of papers [25] - [28], a convergence theorem for Newton's method in Banach spaces is given, which improves the theorems of Kantorovich [4], Lancaster [8] and Ostrowski [16]. The error bounds obtained improve the recent results of Potra [17].

AMS(MOS) Subject Classifications: 65G99, 65J15

Key Words: convergence theorem, Newton's method, Kantorovich's theorem, Lancaster's theorem, Ostrowski's theorem, error estimates, Potra's bounds

Work Unit Number 3 (Numerical Analysis and Scientific Computing)

*Department of Mathematics, Faculty of Science, Ehime University, Matsuyama 790, Japan.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041, and by the Ministry of Education in Japan.


SIGNIFICANCE AND EXPLANATION

To find sharper error bounds for iterative solution of nonlinear equations under assumptions as weak as possible is of basic importance in numerical analysis. This paper gives a convergence theorem for Newton's method in Banach spaces which improves the theorems of Kantorovich [4], Lancaster [8] and Ostrowski [16]. The error bounds obtained improve the recent results of Potra [17].



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.



A CONVERGENCE THEOREM FOR NEWTON'S
METHOD IN BANACH SPACES

Tetsuro Yamamoto*

1. Introduction

There is much literature concerning convergence and error estimates for Newton's method in Banach spaces. In a series of papers [25] - [28], we examined the error bounds which have been obtained by many authors (Dennis [1], Tapia [24], Rall-Tapia [20], Ostrowski [15], [16], Gragg-Tapia [3], Miel [9] - [11], Potra-Pták [18], Moret [12]) under the assumptions of the Kantorovich theorem, and compared them with the Kantorovich bounds. As the result, we concluded [28] that their results follow from the Kantorovich theorem so that, under the Kantorovich assumptions, the Kantorovich theorem still give the best upper bounds for the Newton method.

In this paper, we are interested in improving the assumptions of the Kantorovich theorem and the assertions of the Ostrowski theorem [16; Theorem 38.1]. We shall first state both theorems and several lemmas in §2. Next, in §3, we shall present a convergence theorem which improves both theorems. It will also be shown that results improve the error bounds of Lancaster [8], Kornstaedt [7] and Potra [17]. Finally, in §4, we shall show that Ostrowski's other theorem [16; Theorem 38.2] can be derived by our approach.

2. Preliminaries

Let X and Y be Banach spaces and consider an operator $F : D \subseteq X \rightarrow Y$. If F is Fréchet differentiable in an open convex set $D_0 \subseteq D$, then the Newton method for solving the equation

*Department of Mathematics, Faculty of Science, Ehime University, Matsuyama 790, Japan.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and by the Ministry of Education in Japan.

$$F(x) = 0 \quad (2.1)$$

is defined by

$$x_{n+1} = x_n - F'(x_n)^{-1}F(x_n), \quad n \geq 0, \quad (2.2)$$

provided that $F'(x_n)^{-1} \in L(Y, X)$ exists at each step, where $L(Y, X)$ denotes the Banach space of bounded linear operators of Y into X . Sufficient conditions for convergence of the iterates (2.2), error estimates and existence and uniqueness regions of solutions are given by the famous Kantorovich theorem:

Theorem 2.1 (Kantorovich [4], [5] and Kantorovich-Akilov [6]). Let $F : D \subseteq X \rightarrow Y$ be Frécht differentiable in an open convex set $D_0 \subseteq D$. Assume that for some $x_0 \in D_0$, $F'(x_0)$ is invertible and that

$$\|F'(x_0)^{-1}(F'(x) - F'(y))\| \leq K\|x - y\|, \quad K > 0, \quad x, y \in D_0,$$

$$\|F'(x_0)^{-1}F(x_0)\| \leq \eta, \quad \eta > 0,$$

$$h = K\eta \leq \frac{1}{2}$$

and

$$\bar{S}(x_0, t^*) = \{x \in X \mid \|x - x_0\| \leq t^* = \frac{1 - \sqrt{1 - 2h}}{K}\} \subseteq D_0.$$

Then:

- (i) The iterates (2.2) are well-defined, lie in the open ball $S(x_0, t^*) = \{x \in X \mid \|x - x_0\| < t^*\}$ and converge to a solution x^* of the equation (2.1).
- (ii) The solution x^* is unique in $S(x_0, t^{**}) \cap D_0$ if $2h < 1$ and in $\bar{S}(x_0, t^{**})$ if $2h = 1$, where $t^{**} = (1 + \sqrt{1 - 2h})/K$.
- (iii) Error estimates

$$\|x^* - x_n\| \leq \frac{2\eta_n}{1 + \sqrt{1 - 2h_n}} \leq 2^{1-n}(2h)^{2^n-1}\eta, \quad n \geq 0, \quad (2.3)$$

hold, where η_n and h_n are defined by the recurrence relations

$$B_0 = 1, \quad \eta_0 = \eta, \quad h_0 = h = K\eta,$$

$$B_n = \frac{B_{n-1}}{1 - h_{n-1}}, \quad \eta_n = \frac{h_{n-1}\eta_{n-1}}{2(1 - h_{n-1})}, \quad h_n = KB_n\eta_n, \quad n \geq 1. \quad (2.4)$$

(iii)' Put $f(t) = \frac{1}{2} Kt^2 - t + \eta$ and define the sequence $\{t_n\}$ by

$$t_0 = 0, \quad t_{n+1} = t_n - f(t_n)/f'(t_n), \quad n \geq 0.$$

Then

$$\|x_{n+1} - x_n\| \leq t_{n+1} - t_n$$

and the error estimate

$$\|x^* - x_n\| \leq t^* - t_n, \quad n \geq 0, \quad (2.5)$$

holds.

The bounds (2.3) are of the form found in [4], while the bound (2.5) is found in [5] and [6]. We should remark here that B_n and η_n are the bounds for $\|F'(x_n)^{-1}F'(x_0)\|$ and $\|x_{n+1} - x_n\|$, respectively. In fact, by induction on n , we have

$$\|F'(x_n)^{-1}F'(x_0)\| = \|I + F'(x_{n-1})^{-1}(F'(x_n) - F'(x_{n-1}))\|^{-1}F'(x_{n-1})^{-1}F'(x_0)\|$$

$$\leq \frac{B_{n-1}}{1 - B_{n-1}K\|x_n - x_{n-1}\|} \leq \frac{B_{n-1}}{1 - h_{n-1}} = B_n,$$

$$x_{n+1} - x_n = -F'(x_n)^{-1}F(x_n)$$

$$= -F'(x_n)^{-1}\{F(x_n) - F(x_{n-1}) - F'(x_{n-1})(x_n - x_{n-1})\}$$

$$= -F'(x_n)^{-1} \int_0^1 \{F'(x_{n-1} + t(x_n - x_{n-1})) - F'(x_{n-1})\}(x_n - x_{n-1}) dt \quad (2.6)$$

and

$$\|x_{n+1} - x_n\| \leq \frac{1}{2} B_n K \|x_n - x_{n-1}\|^2 \leq \frac{1}{2} B_n K \eta_{n-1}^2 = \eta_n.$$

On the other hand, Ostrowski [15], [16] proved the convergence of the Newton method under the assumptions which are slightly different from those of Kantorovich.

Theorem 2.2 (Ostrowski [15; Theorems 38.1 and 40.2]). Let $F: D \subseteq X + Y$ and D^0 be the interior of D . Assume that for some $x_0 \in D^0$, $F'(x_0)$ and $F'(x_0)^{-1}$ exist. Let $\varphi \geq 0$, $\alpha = 1 + \cosh \varphi$, $\rho = e^{-\varphi} \|x_1 - x_0\|$ and $\sigma = \alpha \|F(x_0)\| \cdot \|F'(x_0)^{-1}\|^2$. Consider the line segment $L = \{tx_0 + (1-t)x_1 \mid 0 \leq t \leq 1\}$ and the closed ball $\bar{S} = \bar{S}(x_1, \rho)$, and put $C = L \cup \bar{S}$. Assume now that $C \subseteq D^0$, F is Fréchet differentiable on C and

$$\|F'(x) - F'(y)\| \leq \frac{1}{\sigma} \|x - y\|, \quad x, y \in L, \quad x, y \in \bar{S}. \quad (2.7)$$

Then the Newton iterates (2.2) are well-defined, $x_n \in \bar{S}$, $n \geq 1$, and $\{x_n\}$ converges to a solution $x^* \in \bar{S}$ of (2.1), which is unique in C . Furthermore, the following inequalities hold:

$$\|x^* - x_n\| \leq e^{-2^{n-1}\varphi} \frac{\sinh \varphi}{\sinh 2^{n-1}\varphi} \|x_1 - x_0\| \quad (2.8)$$

$$\leq 2^{1-n} \|x_1 - x_0\|,$$

$$\|x^* - x_{n+1}\| \leq e^{-2^n \varphi} \|x_{n+1} - x_n\|, \quad n \geq 0. \quad (2.9)$$

In [28], we derived (2.8) and (2.9) under the assumptions of Theorem 2.1 and showed that they do not improve Gragg-Tapia's bounds. Furthermore, we proved that Moret's bounds, which also follow from Theorem 2.1, are sharper than those of Gragg-Tapia, Potra-Pták and Miel. The argument in [28] also works under the assumptions of an affine invariant version of Theorem 2.2, which are weaker than those of Theorem 2.1 with $\eta = \|x_1 - x_0\|$, provided that $x_0 \neq x_1$. Therefore, on the basis of results obtained in [28], we can improve Theorems 2.1 and 2.2.

Before giving an improved version of both theorems, we state several lemmas. In the following, without loss of generality, we assume that $F(x_0) \neq 0$. This assumption will be kept throughout this paper.

Lemma 2.1. Let $F: D \subseteq X + Y$ and D^0 be the interior of D . Assume that for some $x_0 \in D^0$, $F'(x_0)$ and $F'(x_0)^{-1}$ exist and $F(x_0) \neq 0$. Let $\varphi \geq 0$, $\alpha = 1 + \cosh \varphi$,

$\eta = \|x_1 - x_0\|$ and $\rho = e^{-\alpha}\eta$. Define the sets L , \bar{S} and C as in Theorem 2.2. Furthermore, assume that $C \subseteq D^0$, F is Fréchet differentiable on C and, for some $K > 0$,

$$\|F'(x_0)^{-1}(F'(x) - F'(y))\| \leq K\|x - y\|, \quad x, y \in L, \quad x, y \in \bar{S}, \quad (2.10)$$

and

$$\alpha h \equiv \alpha K \eta \leq 1.$$

Then the iterates (2.2) are well-defined, $x_n \in S = S(x_1, \rho)$ (open ball), $n \geq 1$ and $\|x_{n+1} - x_n\| \leq t_{n+1} - t_n$, $n \geq 0$, where $\{t_n\}$ is the majorizing sequence defined in Theorem 2.1. Therefore, the sequence $\{x_n\}$ converges to a solution $x^* \in \bar{S}$ of (2.1) and $\|x^* - x_n\| \leq t^* - t_n$.

Proof. By the assumption $\alpha h \leq 1$, we have $2h \leq 1$ since $\alpha \geq 2$. Therefore, the majorant theory of Kantorovich can be applied to the sequence $\{x_n\}$, by noting that the condition (2.10) holds and that

$$\begin{aligned} \|x_{n+1} - x_1\| &\leq \|x_{n+1} - x_n\| + \|x_n - x_1\| \\ &\leq (t_{n+1} - t_n) + (t_n - t_1) = t_{n+1} - t_1 \\ &< t^* - t_1 = \frac{1 - h - \sqrt{1 - 2h}}{K} \\ &\leq e^{-\alpha}\eta = \rho, \end{aligned} \quad (2.11)$$

where equality holds in (2.11) if and only if $\alpha h = 1$.

Q.E.D.

Lemma 2.2. Under the assumptions of Lemma 2.1, define the sequences $\{t_n\}$, $\{B_n\}$, $\{\eta_n\}$ and $\{h_n\}$ as in Theorem 2.1. Then

$$\begin{aligned} t_{n+1} - t_n &= \eta_n, \\ t^* - t_n &= \frac{1 - \sqrt{1 - 2h_n}}{KB_n} = \frac{2\eta_n}{1 + \sqrt{1 - 2h_n}} \end{aligned}$$

and

$$t^{**} - t_n = \frac{1 + \sqrt{1 - 2h_n}}{KB_n}, \quad n \geq 0.$$

That is, $t^* - t_n$ and $t^{**} - t_n$ are the solutions of the equation

$$\frac{1}{2} KB_n t^2 - t + \eta_n = 0.$$

Proof. The same proof as in [28] works under the assumptions of Lemma 2.1. Q.E.D.

Lemma 2.3. Under the assumptions of Lemma 2.2, we have for $n \geq 1$

$$\begin{aligned} \text{(i)} \quad B_n^{-1} &= 1 - Kt_n = \sqrt{1 - 2h + (K\eta_{n-1})^2} \\ \text{(ii)} \quad \frac{1}{2} KB_n &= \frac{\nabla t_{n+1}}{(\nabla t_n)^2}, \\ \text{(iii)} \quad \frac{KB_n}{1 + \sqrt{1 - 2h_n}} &= \frac{t^* - t_n}{(\nabla t_n)^2}, \end{aligned}$$

where ∇ denotes the backward difference operator.

Proof. See the proof of Proposition A.3 in [28].

Q.E.D.

Lemma 2.4. Under the assumptions of Lemma 2.2, let $\theta = t^*/t^{**} = (1 - \sqrt{1 - 2h})/(1 + \sqrt{1 - 2h})$. Then we have for $n \geq 0$

$$\begin{aligned} \text{(i)} \quad t^* - t_n &= \begin{cases} \frac{2}{K} \sqrt{1 - 2h} \frac{\theta^{2^n}}{1 - \theta^{2^n}} & (2h < 1) \\ 2^{1-n} \eta & (2h = 1) \end{cases}, \\ \text{(ii)} \quad \frac{t^* - t_{n+1}}{\nabla t_{n+1}} &= \theta^{2^n}. \end{aligned}$$

Proof. See the proofs of Proposition A.1 and Proposition A.4 (ii)

in [28].

Q.E.D.

Lemma 2.5. Under the notation and assumptions of Lemma 2.4, we have for $n \geq 0$

$$(i) \quad t^* - t_n \leq \begin{cases} e^{-2^{n-1}\varphi} \frac{\sinh \varphi}{\sinh 2^{n-1}\varphi} \eta & (\varphi > 0) \\ 2^{1-n}\eta = \lim_{\varphi \rightarrow +0} (e^{-2^{n-1}\varphi} \frac{\sinh \varphi}{\sinh 2^{n-1}\varphi} \eta) & (\varphi = 0) \end{cases},$$

$$(ii) \quad \theta 2^n \leq e^{-2^n \varphi}.$$

The equalities hold in (i) and (ii) if and only if $\alpha h = 1$.

Proof. Take $\varphi^* \geq 0$ such that $\alpha^* h \equiv (1 + \cosh \varphi^*)h = 1$. Then, by Proposition A.4 in [28], we have that the equalities hold in (i) and (ii). Therefore Lemma 2.5 follows for every $\varphi \in [0, \varphi^*]$, since the right hand-sides of (i) and (ii) are monotone decreasing with respect to φ . Q.E.D.

We end this section by proving the following lemma.

Lemma 2.6. Under the assumptions of Lemma 2.1, define the sequence $\{B_n\}$ as in Theorem 2.1. If, for some n , there exists a constant $M_n > 0$ such that

$$\|x^* - x_{n+1}\| \leq \frac{1}{2} M_n \|x^* - x_n\|^2$$

and $M_n \leq KB_n$. Then

$$\|x^* - x_n\| \leq \sigma_n^* \equiv \frac{2d_n}{1 + \sqrt{1 - 2M_n d_n}},$$

where $d_n = \|x_{n+1} - x_n\|$.

Proof. Without loss of generality, we may assume that $d_n \neq 0$, that is $x^* \neq x_n$.

Then, by assumptions, we have

$$\|x^* - x_n\| - d_n \leq \frac{1}{2} M_n \|x^* - x_n\|^2 \leq \frac{1}{2} KB_n \|x^* - x_n\|^2.$$

Hence, if we put

$$\phi_n(t) = \frac{1}{2} M_n t^2 - t + d_n \quad \text{and} \quad \tilde{\phi}_n(t) = \frac{1}{2} KB_n t^2 - t + d_n,$$

then

$$\tilde{\phi}_n(\|x^* - x_n\|) \geq \phi_n(\|x^* - x_n\|) \geq 0$$

and

$$\tilde{\phi}_n(t) > \phi_n(t) \text{ for } t > 0 \text{ or } \tilde{\phi}_n(t) \equiv \phi_n(t).$$

By Lemmas 2.1 and 2.2, we have

$$\|x^* - x_n\| \leq t^* - t_n$$

and $t^* - t_n$, $t^{**} - t_n$ are two solutions of the equation $\psi_n(t) = \frac{1}{2} KB_n t^2 - t + \eta_n = 0$.

Furthermore we have $\psi_n(t) \geq \tilde{\phi}_n(t)$. Therefore $\phi_n(t)$ and $\tilde{\phi}_n(t)$ have positive solutions

σ_n^* , σ_n^{**} and $\tilde{\sigma}_n^*$, $\tilde{\sigma}_n^{**}$ respectively such that

$$\sigma_n^* < \tilde{\sigma}_n^* \leq t^* - t_n \leq t^{**} - t_n \leq \tilde{\sigma}_n^{**} < \sigma_n^{**} \text{ if } M_n < KB_n,$$

$$\sigma_n^* = \tilde{\sigma}_n^* < t^* - t_n \leq t^{**} - t_n < \tilde{\sigma}_n^{**} = \sigma_n^{**} \text{ if } M_n = KB_n \text{ and } d_n < \eta_n$$

and

$$\sigma_n^* = \tilde{\sigma}_n^* = t^* - t_n \leq t^{**} - t_n = \tilde{\sigma}_n^{**} = \sigma_n^{**} \text{ if } M_n = KB_n \text{ and } d_n = \eta_n.$$

In any case we have

$$\|x^* - x_n\| \leq \sigma_n^*,$$

since $\phi_n(\|x^* - x_n\|) \geq 0$ implies $\|x^* - x_n\| \leq \sigma_n^*$ or $\|x^* - x_n\| \geq \sigma_n^{**}$

and the latter case can be excluded.

Q.E.D.

3. Results

We are now in a position to prove the following theorem.

Theorem 3.1. Under the assumptions of Lemma 2.1, the following results hold:

(i) The iterates (2.2) are well-defined, the sequence $\{x_n\}$, $n \geq 1$ remains in an open ball $S = S(x_1, \rho)$ and converges to a solution $x^* \in \bar{S}$ of the equation (2.1).

(ii) The solution is unique in C .

(iii) Let

$$K_n = \sup_{\substack{x, y \in S \\ x \neq y}} \frac{|F'(x_n)^{-1}(F'(x) - F'(y))|}{|x - y|}, \quad n \geq 1,$$

$$K_0 = K$$

and put $d_n = |x_{n+1} - x_n|$. Then the following error estimates hold.

(a) A posteriori error estimates:

$$|x^* - x_n| \leq \frac{2d_n}{1 + \sqrt{1 - 2K_n d_n}} \quad (n \geq 0) \quad (3.1)$$

$$\leq \begin{cases} t^* = \frac{2\eta}{1 + \sqrt{1 - 2h}} & (n = 0) \\ \frac{2d_n}{1 + \sqrt{1 - 2K[1 - K(|x_n - x_1| + d_0)]^{-1}d_n}} & (n \geq 1) \end{cases} \quad (3.2)$$

$$\leq \frac{2d_n}{1 + \sqrt{1 - 2K(1 - Kt_n)^{-1}d_n}} \quad (n \geq 0) \quad (3.3)$$

$$\leq \frac{t^* - t_n}{\sqrt{t_{n+1}}} d_n \quad (n \geq 0)$$

$$\leq \frac{t^* - t_n}{(\sqrt{t_n})^2} d_{n-1}^2 \quad (n \geq 1) \quad (\text{Miel [10]})$$

$$\leq \frac{Kd_{n-1}^2}{\sqrt{1 - 2h + (Kd_{n-1})^2} + \sqrt{1 - 2h}} \quad (n \geq 1) \quad (\text{Potra-Ptáček [18]})$$

$$\leq \frac{t^* - t_n}{\sqrt{t_n}} d_{n-1} \quad (n \geq 1) \quad (\text{Miel [10]})$$

$$= \theta^{2^{n-1}} d_{n-1} \quad (n \geq 1) \quad (\text{Gragg-Tapia [3]})$$

$$\leq e^{-2^{n-1}\varphi} d_{n-1} \quad (n \geq 1) \quad (\text{Ostrowski [16]}). \quad (3.4)$$

The equality holds in (3.4) if and only if $ah = 1$.

(b) A priori error estimates:

$$\|x^* - x_n\| \leq t^* - t_n \quad (n \geq 0) \quad (\text{Kantorovich [5], [6]})$$

$$= \begin{cases} \frac{2}{K} \sqrt{1-2h} \frac{\theta^{2^n}}{1-\theta^{2^n}} & (2h < 1) \\ 2^{1-n} \eta & (2h = 1) \quad (n \geq 0) \quad (\text{Gragg-Tapia [3]}) \end{cases}$$

$$\leq \begin{cases} e^{-2^{n-1}\varphi} \frac{\sinh \varphi}{\sinh 2^{n-1}\varphi} \eta & (\varphi > 0) \\ 2^{1-n} \eta & (\varphi = 0) \quad (n \geq 0). \quad (\text{Ostrowski [16]}) \end{cases} \quad (3.5)$$

The equality holds in (3.5) if and only if $ah = 1$.

(iv) If F is Fréchet differentiable in an open convex set D_0 such that $D^0 \supseteq D_0 \supset C$ and if $F'(x)$ satisfies the Lipschitz condition in D_0 with the Lipschitz constant K , then the solution x^* is unique in

$$\tilde{S} = \begin{cases} S(x_0, t^{**}) \cap D_0 & \text{if } 2h < 1 \\ \bar{S}(x_0, t^{**}) \cap D_0 & \text{if } 2h = 1. \end{cases}$$

Furthermore, (3.2) may be replaced by the sharper bound

$$\|x^* - x_n\| \leq \frac{2d_n}{1 + \sqrt{1 - 2K(1 - K\Delta_n)^{-1}d_n}} \quad (n \geq 0) \quad (\text{Moret [12]}) \quad (3.6)$$

where $\Delta_n = \|x_n - x_0\|$.

Proof. (i) was proved in Lemma 2.1. To prove (ii), let \tilde{x}^* be a solution in \bar{S} . Then we have

$$\begin{aligned}\tilde{x}^* - x_{n+1} &= \tilde{x}^* - x_n + F'(x_n)^{-1}F(x_n) \\ &= -F'(x_n)^{-1}\{F(\tilde{x}^*) - F(x_n) - F'(x_n)(\tilde{x}^* - x_n)\} \\ &= -F'(x_n)^{-1}F'(x_0) \int_0^1 F'(x_0)^{-1}\{F'(x_n + t(\tilde{x}^* - x_n)) - F'(x_n)\}(\tilde{x}^* - x_n)dt\end{aligned}$$

and

$$x_n + t(\tilde{x}^* - x_n), \quad x_n \in \bar{S}, \quad n \geq 1.$$

Hence, by (2.10) and Lemma 2.3 (ii) we have

$$\|\tilde{x}^* - x_{n+1}\| \leq \frac{1}{2} B_n K \|\tilde{x}^* - x_n\|^2 = \frac{\nabla t_{n+1}}{(\nabla t_n)^2} \|\tilde{x}^* - x_n\|^2$$

so that

$$\frac{\|\tilde{x}^* - x_{n+1}\|}{\nabla t_{n+1}} \leq \left(\frac{\|\tilde{x}^* - x_n\|}{\nabla t_n} \right)^2 \leq \dots \leq \left(\frac{\|\tilde{x}^* - x_1\|}{\nabla t_1} \right)^{2^n} \leq \left(\frac{\rho}{\eta} \right)^{2^n} \leq 1.$$

This implies

$$\|\tilde{x}^* - x_{n+1}\| \leq \nabla t_{n+1} \rightarrow 0$$

as $n \rightarrow \infty$. Thus we obtain $\tilde{x}^* = \lim_{n \rightarrow \infty} x_n = x^*$. Next, we shall show that there is no solution in $C \setminus \bar{S}$, provided that the set $C \setminus \bar{S}$ is not empty. To show this, let \tilde{x}^* be a solution in $C \setminus \bar{S}$. Then we have

$$e^{-\varphi} \eta = \rho < \|\tilde{x}^* - x_1\| \leq \frac{1}{2} K \|\tilde{x}^* - x_0\|^2 < \frac{1}{2} K (\eta - \rho)^2 = \frac{1}{2} K (1 - e^{-\varphi})^2 \eta^2$$

so that

$$1 < K (\cosh \varphi - 1) \eta = K(a - 2) \eta < 1 - 2h,$$

which is a contradiction. This proves the uniqueness of the solution in C . To prove (iii), we first observe that

$$x^* - x_{n+1} = -F'(x_n) \int_0^1 \{F'(x_n + t(x^* - x_n)) - F'(x_n)\} (x^* - x_n) dt$$

$$x_n + t(x^* - x_n), \quad x_n \in \bar{S}, \quad n \geq 1,$$

$$F'(x_n)^{-1} = \{I + F'(x_0)^{-1}(F'(x_n) - F'(x_1)) + F'(x_0)^{-1}(F'(x_1) - F'(x_0))\}^{-1} F'(x_0)^{-1}$$

and

$$x_n, x_1 \in \bar{S}, \quad n \geq 1, \quad x_1, x_0 \in L.$$

Hence we have from Lemma 2.3 (i)

$$\|F'(x_n)^{-1}F'(x_0)\| \leq B_n' \equiv \frac{1}{1 - K(\|x_n - x_1\| + d_0)} \quad (n \geq 1) \quad (3.7)$$

$$\leq \frac{1}{1 - Kt_n} = B_n$$

and

$$\|x^* - x_{n+1}\| \leq \frac{1}{2} K_n \|x^* - x_n\|^2 \leq \frac{1}{2} K B_n' \|x^* - x_n\|^2 \leq \frac{1}{2} K B_n \|x^* - x_n\|^2.$$

Therefore Lemma 2.6 can be applied to obtain the bounds (3.1) - (3.3) for $n \geq 1$. Observe

that if $n = 0$, they reduce to the Kantorovich bound $t^* = 2\eta/(1 + \sqrt{1 - 2h})$. The other

part of (iii) follows from Lemmas 2.2 - 2.5. This proves (iii). Finally we shall prove

(iv). Let F be Fréchet differentiable in an open convex set D_0 such that

$D^0 \supseteq D_0 \supset C$ and $F'(x)$ satisfy the Lipschitz condition (2.10) in D_0 . Then (3.7) may be replaced by the sharper estimate

$$\|F'(x_n)^{-1}F'(x_0)\| \leq \frac{1}{1 - K\|x_n - x_0\|}, \quad n \geq 0.$$

Therefore, Lemma 2.6 can again be applied to replace (3.2) by (3.6). To prove the

uniqueness of solution in \tilde{S} , let \tilde{x}^* be a solution in \tilde{S} . Then there exists a

nonnegative constant r such that $r < 1$ if $2h < 1$, $r \leq 1$ if $2h = 1$ and

$\|\tilde{x}^* - x_0\| \leq rt^{**}$. By induction on n , we can show that

$$\|\tilde{x}^* - x_n\| \leq r^{2^n} (t^{**} - t_n), \quad n \geq 0. \quad (3.8)$$

In fact, we have under our assumptions

$$\begin{aligned} \|\tilde{x}^* - x_{n+1}\| &\leq \frac{1}{2} K B_n \|x^* - x_n\|^2 \\ &\leq \frac{1}{2} K B_n \{r^{2^n} (t^{**} - t_n)\}^2 \\ &= r^{2^{n+1}} (t^{**} - t_n - \eta_n) \\ &= r^{2^{n+1}} (t^{**} - t_{n+1}), \end{aligned}$$

where we have used the induction hypothesis and Lemma 2.2. This proves (3.8), from which we obtain

$$\|\tilde{x}^* - x_n\| \rightarrow 0$$

as $n \rightarrow \infty$, since $r^{2^n} \rightarrow 0$ if $2h < 1$ and $t^{**} - t_n = t^* - t_n \rightarrow 0$ if $2h = 1$. Hence we have $\tilde{x}^* = \lim_{n \rightarrow \infty} x_n = x^*$, which implies the uniqueness of solution in \tilde{S} . Q.E.D.

Assumptions similar to those of Theorem 2.2 or Lemma 2.1 were also adopted by Lancaster [8], and later by Schmidt [22], [23] for the generalized secant method which includes the Newton method as a special case. Lancaster's assumptions correspond to the case $\varphi = 0$ in Lemma 2.1, while Schmidt's correspond to the case where φ is chosen so that $ah = 1$, in which case we have $\rho = e^{-\varphi} \eta = t^* - \eta$. In the following we shall improve Lancaster's result. (In the case of the Newton method, Schmidt's upper bound reduces to the Kantorovich bound $t^* - t_n$. Also see [28].)

Corollary 3.1.1. Let $F: D \subseteq X \rightarrow Y$ and D^0 be the interior of D . Assume that for some $x_0 \in D^0$, $F'(x_0)$ and $F'(x_0)^{-1}$ exist, $F'(x_0) \neq 0$ and F is Fréchet differentiable on $\bar{S}_0 = \bar{S}(x_1, \|x_1 - x_0\|)$. Furthermore, put

$$L_0 = \sup_{\substack{x, y \in \bar{S}_0 \\ x \neq y}} \frac{\|F'(x_0)^{-1}(F'(x) - F'(y))\|}{\|x - y\|}.$$

If $2L_0\|x_1 - x_0\| \leq 1$, then the Newton process (2.2) generates a sequence $\{x_n\} \subset \bar{S}_0$ which converges to the unique solution x^* of (2.1) in \bar{S}_0 . If we put

$$L_n = \sup_{\substack{x, y \in \bar{S}_0 \\ x \neq y}} \frac{\|F'(x_n)^{-1}(F'(x) - F'(y))\|}{\|x - y\|}, \quad n \geq 1$$

and

$$d_n = \|x_{n+1} - x_n\|, \quad n \geq 0,$$

then the following error estimates hold:

$$\|x^* - x_n\| \leq \delta_n \equiv \frac{2d_n}{1 + \sqrt{1 - 2L_n d_n}} \quad (n \geq 0)$$

$$\leq \frac{L_n d_{n-1}^2}{1 + \sqrt{1 - (L_n d_{n-1})^2}} \quad (n \geq 1)$$

$$\leq \frac{L_{n-1} d_{n-1}^2}{1 - L_{n-1} d_{n-1} + \sqrt{1 - 2L_{n-1} d_{n-1}}} \quad (n \geq 1) \quad (3.9)$$

$$\leq \frac{L_{n-1} d_{n-1}^2}{1 - L_{n-1} d_{n-1}} \quad (n \geq 1). \quad (3.10)$$

Proof. Put $\varphi = 0$ in Theorem 3.1. Then Corollary 3.1.1 follows from Theorem 3.1 by noting that

$$d_n \leq \frac{1}{2} L_n d_{n-1}^2 \quad \text{and} \quad L_n \leq \frac{L_{n-1}}{1 - L_{n-1} d_{n-1}}, \quad n \geq 1. \quad \text{Q.E.D.}$$

The bound (3.10) is due to Lancaster [8] and (3.9) is what Potra cited in his recent paper [17] as Kornstaedt's bound [7].

Corollary 3.1.2. Under the notation and assumptions of Corollary 3.1.1, we have

$$\begin{aligned} \|x^* - x_n\| &\leq \delta_n \leq \frac{2d_n}{1 + \sqrt{1 - 2L_0(1 - L_0\Delta_n)^{-1}d_n}} \quad (n \geq 0) \\ &\leq \frac{2\|F'(x_0)^{-1}F(x_n)\|}{1 - L_0\Delta_n + \sqrt{(1 - L_0\Delta_n)^2 - 2L_0\|F'(x_0)^{-1}F(x_n)\|}} \quad (n \geq 0) \end{aligned} \quad (3.11)$$

$$\leq \frac{L_0 d_{n-1}^2}{1 - L_0\Delta_n + \sqrt{(1 - L_0\Delta_n)^2 - (L_0 d_{n-1})^2}} \quad (n \geq 1) \quad (3.12)$$

where $\Delta_n = \|x_n - x_0\|$.

Proof. It is easy to see that

$$L_n \leq \frac{L_0}{1 - L_0\Delta_n}$$

and

$$\begin{aligned} d_n &\leq \|F'(x_n)^{-1}F'(x_0)\| \cdot \|F'(x_0)^{-1}F(x_n)\| \\ &\leq \frac{1}{1 - L_0\Delta_n} \|F'(x_0)^{-1}F(x_n)\|. \end{aligned}$$

Furthermore, we have

$$\begin{aligned} F'(x_0)^{-1}F(x_n) &= F'(x_0)^{-1}\{F(x_n) - F(x_{n-1}) - F'(x_{n-1})(x_n - x_{n-1})\} \\ &= F'(x_0)^{-1} \int_0^1 \{F'(x_{n-1} + t(x_n - x_{n-1})) - F'(x_{n-1})\}(x_n - x_{n-1}) dt \end{aligned}$$

so that

$$\|F'(x_0)^{-1}F(x_n)\| \leq \frac{1}{2} L_0 d_{n-1}^2, \quad n \geq 1.$$

Therefore, Corollary 3.1.2 follows from Corollary 3.1.1.

Q.E.D.

Remark 3.1. The bounds (3.11) and (3.12) were recently obtained by Potra [17] under the assumptions of Theorem 2.1 as $\beta_5(\Delta_n, \|F'(x_0)^{-1}F(x_n)\|)$ and $\beta_4(\Delta_n, d_{n-1})$ respectively in his notation.

We can further improve the bound δ_n obtained in Corollary 3.1.1.

Corollary 3.1.3. Under the assumptions of Corollary 3.1.1, put $M_0 = L_0$ and for

$n \geq 1$

$$\bar{S}_n = \bar{S}(x_n, 2\|F'(x_n)^{-1}F(x_n)\|),$$

$$M_n = \sup_{\substack{x, y \in \bar{S}_n \\ x \neq y}} \frac{\|F'(x_n)^{-1}(F'(x) - F'(y))\|}{\|x - y\|}.$$

Then we have

$$x^* \in \bar{S}_n \subseteq \bar{S}_{n-1} \subseteq \dots \subseteq \bar{S}_0.$$

and

$$\|x^* - x_n\| \leq \epsilon_n \equiv \frac{2d_n}{1 + \sqrt{1 - 2M_n d_n}} \leq \delta_n, \quad n \geq 0.$$

Proof. This immediately follows from Lemma 2.6 by noting that $2d_{n+1} \leq d_n$ and

$$\|x^* - x_{n+1}\| \leq \frac{1}{2} M_n \|x^* - x_n\|^2 \leq \frac{1}{2} L_n \|x^* - x_n\|^2, \quad n \geq 0. \quad \text{Q.E.D.}$$

Remark 3.2. As was remarked in [17], the cost of obtaining K_n , L_n or M_n might be very high. Therefore, in practical computation, it would be better to make use of one of (3.2), (3.3), (3.6) and (3.12). However, Theorem 3.1 and its Corollaries assert that the error bounds which have been obtained by many authors with the use of different techniques can be derived from the majorant theory of Kantorovich and Lemma 2.6, in a unified manner.

Theorem 3.2. Under the assumptions of Lemma 2.1, we have

$$\begin{aligned} d_n &\leq \frac{\sqrt{t_{n+1}}}{(\sqrt{t_n})^2} d_{n-1}^2 \leq \frac{\sqrt{t_{n+1}}}{\sqrt{t_n}} d_{n-1} \\ &\leq \frac{1}{2 \cosh 2^{n-1} \varphi} d_{n-1}, \quad n \geq 1. \end{aligned} \quad (3.13)$$

The equality holds in (3.13) if and only if $\alpha h = 1$.

Proof. It follows from (2.6) and Lemma 2.3 (ii) that

$$d_n \leq \frac{1}{2} K_B d_{n-1}^2 = \frac{v_{t_{n+1}}}{(v_{t_n})^2} d_{n-1}^2 \leq \frac{v_{t_{n+1}}}{v_{t_n}} d_{n-1}. \quad (3.14)$$

Choose $\varphi^* \geq 0$ such that $(1 + \cosh \varphi^*)h = 1$. Then, from Lemmas 2.4 and 2.5, we have

$$v_{t_{n+1}} = e^{2^n \varphi^* (t^* - t_{n+1})} = \begin{cases} \frac{\sinh \varphi^*}{\sinh 2^n \varphi^*} & (\varphi^* > 0) \\ 2^{-n} & (\varphi^* = 0) \end{cases}$$

Hence

$$\begin{aligned} \frac{v_{t_{n+1}}}{v_{t_n}} &= \begin{cases} \frac{\sinh 2^{n-1} \varphi^*}{\sinh 2^n \varphi^*} & (\varphi^* > 0) \\ \frac{1}{2} & (\varphi^* = 0) \end{cases} \\ &= \frac{1}{2 \cosh 2^{n-1} \varphi^*} \leq \frac{1}{2 \cosh 2^{n-1} \varphi} \end{aligned} \quad (3.15)$$

for every $\varphi \in [0, \varphi^*]$. The equality holds in (3.15) if and only if $\varphi = \varphi^*$. This,

together with (3.14), proves Theorem 3.2.

Q.E.D.

Remark 3.3. The bound (3.13) is of the form found in Ostrowski [16; Theorem 38.2].

4. Observation

Under the assumptions of Theorem 2.2, Ostrowski proved that

$$\begin{aligned} \frac{|F(x_{n+m})|}{|F(x_n)|} &\leq \begin{cases} \left(\frac{\sinh 2^{n-1} \varphi}{\sinh 2^{n+m-1} \varphi} \right)^2 & (\varphi > 0) \\ \frac{1}{2^{2m}} = \lim_{\varphi \rightarrow 0} \left(\frac{\sinh 2^{n-1} \varphi}{\sinh 2^{n+m-1} \varphi} \right)^2 & (\varphi = 0) \end{cases} \\ &\leq \frac{1}{2^{2m}}, \quad n \geq 0, \quad m \geq 0, \end{aligned} \quad (4.1)$$

provided that $F(x_n) \neq 0$. By our approach, we can easily derive his estimates. Let

$\bar{K} = \sigma^{-1}$, $\bar{B} = \|F'(x_0)^{-1}\|$, $\bar{\eta} = \bar{B}\|F(x_0)\|$, $\bar{f}(t) = \frac{1}{2}\bar{K}\bar{B}t^2 - t + \bar{\eta}$. Furthermore, we define the sequences $\{\bar{B}_n\}$ and $\{\bar{\eta}_n\}$ by

$$\bar{B}_0 = \bar{B}, \quad \bar{\eta}_0 = \bar{\eta}, \quad \bar{h}_0 = \bar{K}\bar{B}_0\bar{\eta}_0,$$

$$\bar{B}_n = \frac{\bar{B}_{n-1}}{1 - \bar{h}_{n-1}}, \quad \bar{\eta}_n = \frac{\bar{h}_{n-1}\bar{\eta}_{n-1}}{2(1 - \bar{h}_{n-1})}, \quad \bar{h}_n = \bar{K}\bar{B}_n\bar{\eta}_n, \quad n \geq 1.$$

Then it is easy to see that

$$\|F'(x_n)^{-1}\| \leq \bar{B}_n, \quad \|x_{n+1} - x_n\| \leq \bar{\eta}_n$$

and

$$\begin{aligned} F(x_{n+1}) &= F(x_n) + \int_0^1 F'(x_n + t(x_{n+1} - x_n))(x_{n+1} - x_n) dt \\ &= \int_0^1 \{F'(x_n) - F'(x_n + t(x_{n+1} - x_n))\} F'(x_n)^{-1} F(x_n) dt. \end{aligned}$$

Hence we have

$$\begin{aligned} \|F(x_{n+1})\| &\leq \frac{1}{2} \bar{K} \|x_{n+1} - x_n\| \|F'(x_n)^{-1}\| \|F(x_n)\| \\ &\leq \frac{1}{2} \bar{K} \bar{\eta}_n \bar{B}_n \|F(x_n)\| = \left(\frac{\bar{\eta}_n}{\bar{\eta}_{n-1}} \right)^2 \|F(x_n)\|. \end{aligned} \quad (4.2)$$

Define the sequence $\{\bar{t}_n\}$ by

$$\bar{t}_0 = 0, \quad \bar{t}_{n+1} = \bar{t}_n - \bar{f}(\bar{t}_n) / \bar{f}'(\bar{t}_n), \quad n \geq 0.$$

Then, by Lemmas 2.2, 2.4 and 2.5, we have

$$\bar{\eta}_n = \bar{\epsilon}_{n+1} - \bar{\epsilon}_n = e^{2^n \varphi} (\bar{\epsilon}^* - \bar{\epsilon}_{n+1})$$

$$= \begin{cases} \frac{\sinh \varphi}{\sinh 2^n \varphi} \eta & (\varphi > 0) \\ 2^{-n} \eta & (\varphi = 0) \end{cases} \quad (4.3)$$

since $\alpha \bar{h}_0 = 1$, where $\bar{\epsilon}^* = (1 - \sqrt{1 - 2\bar{h}_0})/\bar{K}$. Therefore, we have from (4.2) and (4.3)

$$\frac{|F(x_{n+1})|}{|F(x_n)|} \leq \begin{cases} \left(\frac{\sinh 2^{n-1} \varphi}{\sinh 2^n \varphi} \right)^2 & (\varphi > 0) \\ \frac{1}{2} & (\varphi = 0) \end{cases}$$

This leads to the estimate (4.1). Therefore, together with (3.13) which holds in our case, we proved the main part of his theorem [16; Theorem 38.2]. The remaining part also follows from our approach.

Finally we remark that the chart for the lower bounds given in [28] is still true under the assumptions of Lemma 2.1 with a slight modification:

$$\begin{aligned} \|x^* - x_n\| &\geq \frac{2d_n}{1 + \sqrt{1 + 2K_n d_n}} \quad (n \geq 0) \\ &\geq \begin{cases} \frac{2d_0}{1 + \sqrt{1 + 2h}} & (n = 0) \\ \frac{2d_n}{1 + \sqrt{1 + 2K\{1 - K(\|x_n - x_1\| + d_0)\}^{-1}d_n}} & (n \geq 1) \end{cases} \quad (4.4) \\ &\geq \frac{2d_n}{1 + \sqrt{1 + 2K(1 - Kt_n)^{-1}d_n}} \quad (n \geq 0) \quad (\text{Miel [11], Schmidt [23]}) \end{aligned}$$

$$\geq \frac{2d_n}{1 + \sqrt{1 + \frac{2Kd_n}{Kd_n + \sqrt{1 - 2h_n + (Kd_n)^2}}}} \quad (n \geq 0) \quad (\text{Potra-Ptak [18]})$$

$$\geq \frac{2d_n}{1 + \sqrt{1 + 2h_n}} \quad (n \geq 0) \quad (\text{Gragg-Tapia [3]}).$$

If the assumptions of Theorem 3.1 (iv) are satisfied, then (4.4) may be replaced by the sharper lower bound

$$\|x^* - x_n\| \geq \frac{2d_n}{1 + \sqrt{1 + 2K(1 - K\Delta_n)^{-1}d_n}} \quad (n \geq 0) \quad (\text{Yamamoto [28]}).$$

We also have

$$\|x^* - x_n\| \geq \epsilon_n \equiv \frac{2d_n}{1 + \sqrt{1 + 2M_n d_n}}, \quad n \geq 0,$$

with the notation and assumptions of Corollary 3.1.3.

Acknowledgements. The author wishes to thank Professor L. B. Rall of the Mathematics Research Center, University of Wisconsin-Madison and Professor R. K. Guy of the University of Calgary for bringing the theorems of Ostrowski and Lancaster respectively to his attention.

References

- [1]. J.E. Dennis, Jr., On the Kantorovich hypothesis for Newton's method, SIAM J. Numer. Anal., 6 (1969), 493-507.
- [2]. P. Deufilhard and G. Heindl, Affine invariant convergence theorems for Newton's method and extensions to related methods, SIAM J. Numer. Anal., 16 (1979), 1-10.
- [3]. W.B. Gragg and R.A. Tapia, Optimal error bounds for the Newton-Kantorovich theorem, SIAM J. Numer. Anal., 11 (1974), 10-13.
- [4]. L.V. Kantorovich, On Newton's method for functional equations, Dokl. Akad. Nauk. SSSR, 59 (1948), 1237-1240.
- [5]. L.V. Kantorovich, On Newton's method, Trudy Mat. Inst. Steklov, 28 (1949), 104-144.
- [6]. L.V. Kantorovich and G.P. Akilov, Functional Analysis in Normed Spaces, Pergamon Press, Oxford, 1964.
- [7]. H.J. Kornstaedt, Funktionalungleichungen und Iterationsverfahren, Aequat. Math., 13 (1975), 21-45.
- [8]. P. Lancaster, Error analysis for the Newton-Raphson Method, Numer. Math., 9 (1966), 55-68.
- [9]. G.J. Miel, The Kantorovich theorem with optimal error bounds, Amer. Math. Monthly, 86 (1979), 212-215.
- [10]. G.J. Miel, Majorizing sequences and error bounds for iterative methods, Math. Comp., 34 (1980), 185-202.
- [11]. G.J. Miel, An updated version of the Kantorovich theorem for Newton's method, Computing, 27 (1981), 237-244.
- [12]. I. Moret, A note on Newton type iterative methods, Computing, 33 (1984), 65-73.
- [13]. J.M. Ortega, The Newton-Kantorovich theorem, Amer. Math. Monthly, 75 (1968), 658-660.
- [14]. J.M. Ortega and W.C. Rheinboldt, Iterative Solution of Nonlinear Equations in Several Variables, Academic Press, New York, 1970.

- [15] A.M. Ostrowski, *La méthode de Newton dans les espaces de Banach*, C.R. Acad. Sci. Paris, 27 (A) (1971), 1251-1253.
- [16] A.M. Ostrowski, *Solution of Equations in Euclidean and Banach Spaces*, Academic Press, New York, 1973.
- [17] F.A. Potra, On the aposteriori error estimates for Newton's method, *Beiträge zur Numerische Mathematik*, 12 (1984), 125-138.
- [18] F.A. Potra and V. Pták, Sharp error bounds for Newton's process, *Numer. Math.*, 34 (1980), 63-72.
- [19] L.B. Rall, *Computational Solution of Nonlinear Operator Equations*, Krieger, Huntington, New York, 1979.
- [20] L.B. Rall and R.A. Tapia, The Kantorovich theorem and error estimates for Newton's method, MRC Technical Summary Report #1043, University of Wisconsin, 1970.
- [21] W.C. Rheinboldt, A unified convergence theory for a class of iterative process, *SIAM J. Numer. Anal.*, 5 (1968), 42-63.
- [22] J.W. Schmidt, Regular-falsi-Verfahren mit Konsistenter Steigung und Majorantenprinzip, *Periodica Math. Hungarica*, 5 (1974), 187-193.
- [23] J.W. Schmidt, Untere Fehlerschranken für Regular-falsi-Verfahren, *Periodica Math. Hungarica*, 9 (1978), 241-247.
- [24] R.A. Tapia, The Kantorovich theorem for Newton's method, *Amer. Math. Monthly*, 78 (1971), 389-392.
- [25] T. Yamamoto, Error bounds for Newton's process derived from the Kantorovich theorem, to appear in *Japan J. Appl. Math.*, 2(1985).
- [26] T. Yamamoto, Error bounds for Newton's iterates derived from the Kantorovich theorem, MRC Technical Summary Report #2843, University of Wisconsin, 1985.
- [27] T. Yamamoto, A unified derivation of several error bounds for Newton's process, *J. Comp. Appl. Math.*, 12 & 13 (1985), 179-191.
- [28] T. Yamamoto, Error bounds for Newton-like methods under Kantorovich type assumptions, MRC Technical Summary Report #2846, University of Wisconsin, 1985.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2879	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A CONVERGENCE THEOREM FOR NEWTON'S METHOD IN BANACH SPACES		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Tetsuro Yamamoto		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Numerical Analysis and Scientific Computing
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE October 1985
		13. NUMBER OF PAGES 22
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) convergence theorem, Newton's method, Kantorovich's theorem, Lancaster's theorem, Ostrowski's theorem, error estimates, Potra's bounds		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) On the basis of the results obtained in a series of papers [25] - [28], a convergence theorem for Newton's method in Banach spaces is given, which improves the theorems of Kantorovich [4], Lancaster [8] and Ostrowski [16]. The error bounds obtained improve the recent results of Potra [17].		

END

FILMED

386

DTIC